



All Content

Search

[Advanced Search](#)

< Previous Article

January 2018 Volume 97, Issue 1, Pages 48–53

Next Article >

 Access this article on
[ScienceDirect](#)

Moderate intravenous sedation for first trimester surgical abortion: a comparison of adverse outcomes between obese and normal-weight women☆☆☆☆

[Gillian Horwitz](#), [Danielle Roncari](#), [Kari P. Braaten](#), [Rie Maurer](#), [Jennifer Fortin](#), [Alisa B. Goldberg](#)

PlumX Metrics

 DOI: <https://doi.org/10.1016/j.contraception.2017.09.001> | CrossMark

Article Info

Article Tools

[PDF \(142 KB\)](#)[Email Article](#)[Add to My Reading List](#)[Export Citation](#)[Create Citation Alert](#)[Cited by in Scopus \(0\)](#)[Request Permissions](#)[Order Reprints](#)

(100 minimum order)

Blood loss at the time of dilation and evacuation at 16 to 22 weeks of gestation in women using low molecular weight heparin: a case series
 Tschann, Mary et al.

[Abstract](#) | [Full Text](#) | **References**

References

Induced abortion among women veterans: data from the ECUUN study
 Schwarz, Eleanor Bimla et al.

1. **Abortion incidence and service availability in the United States, 2014.** *Health Aff (Millwood)*. 2017;36:17–27.
[Crossref](#) | [PubMed](#) | [Scopus \(30\)](#) | [Google Scholar](#)

2. Guttmacher Institute. **Induced abortion in the United States.**
https://www.guttmacher.org/sites/default/files/factsheet/fb_induced_abortion_5.pdf
 Date: 2017
 Date accessed: August 19, 2017
 .
[Google Scholar](#)

3. Henshaw, S.K. **Unintended pregnancy in the United States.** *Fam Plann Perspect*. 1998;30:24–29 ([46]).
[Crossref](#) | [PubMed](#) | [Scopus \(1012\)](#) | [Google Scholar](#)

4. Jones, R.K., Kavanaugh, M.L. **Changes in abortion rates between 2000 and 2008 and lifetime incidence of abortion.** *Obstet Gynecol*. 2011;117:1358–1366.
[Crossref](#) | [PubMed](#) | [Scopus \(111\)](#) | [Google Scholar](#)

5. Flegal, K.M., Carroll, M.D., Kit, B.K., Ogden, C.L. **Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999–2010.** *JAMA*. 2012;307:491–497.
[Crossref](#) | [PubMed](#) | [Scopus \(3085\)](#) | [Google Scholar](#)

6. O'Connell, K., Jones, H.E., Simon, M., Saporta, V., Paul, M., Lichtenberg, E.S. **First-trimester surgical abortion practices: a survey of National Abortion Federation members.** *Contraception*. 2009;79:385–392.
[Abstract](#) | [Full Text](#) | [Full Text PDF](#) | [PubMed](#) | [Scopus \(70\)](#) | [Google Scholar](#)

7. O'Connell, K., Jones, H.E., Lichtenberg, E.S., Paul, M. **Second-trimester surgical abortion practices: a survey of National Abortion Federation members.** *Contraception*. 2008;78:492–499.
[Abstract](#) | [Full Text](#) | [Full Text PDF](#) | [PubMed](#) | [Scopus \(37\)](#) | [Google Scholar](#)

8. American Society of Anesthesiologists. **Continuum of depth of sedation: definition of general anesthesia and levels of sedation/analgesia.**
<http://www.asahq.org/~media/sites/asahq/files/public/resources/standards-guidelines/continuum-of-depth-of-sedation-definition-of-general-anesthesia-and-levels-of-sedation-analgesia.pdf>
 Date: 2014
 Date accessed: July 20, 2017
 .
[Google Scholar](#)

9. Wong, C.Y., Ng, E.H., Ngai, S.W., Ho, P.C. **A randomized, double blind, placebo-controlled study to investigate the use of conscious sedation in conjunction with paracervical block for reducing pain in termination of first trimester pregnancy by suction evacuation.** *Hum Reprod*. 2002;17:1222–1225.
[Crossref](#) | [PubMed](#) | [Google Scholar](#)

10. Rawling, M.J., Wiebe, E.R. **A randomized controlled trial of fentanyl for abortion pain.** *Am J Obstet Gynecol.* 2001;185:103–107.

[Abstract](#) | [Full Text](#) | [Full Text PDF](#) | [PubMed](#) | [Scopus \(37\)](#) | [Google Scholar](#)

11. Allen, R.H., Kumar, D., Fitzmaurice, G., Lifford, K.L., Goldberg, A.B. **Pain management of first-trimester surgical abortion: effects of selection of local anesthesia with and without lorazepam or intravenous sedation.** *Contraception.* 2006;74:407–413.

[Abstract](#) | [Full Text](#) | [Full Text PDF](#) | [PubMed](#) | [Scopus \(28\)](#) | [Google Scholar](#)

12. Allen, R.H., Fitzmaurice, G., Lifford, K.L., Lasic, M., Goldberg, A.B. **Oral compared with intravenous sedation for first-trimester surgical abortion.** *Obstet Gynecol.* 2009;113:276–283.

[Crossref](#) | [PubMed](#) | [Scopus \(13\)](#) | [Google Scholar](#)

13. Allen, R.H., Fortin, J., Bartz, D., Goldberg, A.B., Clark, M.A. **Women's Preferences for pain control during first-trimester surgical abortion: a qualitative study.** *Contraception.* 2012;85:413–418.

[Abstract](#) | [Full Text](#) | [Full Text PDF](#) | [PubMed](#) | [Scopus \(9\)](#) | [Google Scholar](#)

14. National Institutes of Health. **Classification of overweight and obesity by BMI, waist circumference, and associated disease risks.**

https://www.nhlbi.nih.gov/health/educational/lose_wt/BMI/bmi_dis.htm

Date accessed: July 20, 2017

[Google Scholar](#)

15. Strum, R. **Increases in morbid obesity in the USA: 2000-2005.** *Public Health.* 2007;121:492–496.

[Abstract](#) | [Full Text](#) | [Full Text PDF](#) | [PubMed](#) | [Scopus \(440\)](#) | [Google Scholar](#)

16. Segal, L.M., Martin, A., Rayburn, J. **The state of obesity: better policies for a healthier America.**

<http://stateofobesity.org/files/stateofobesity2016.pdf>

Date: 2016

Date accessed: July 20, 2017

[Google Scholar](#)

17. Karamnov, S., Sarkisian, N., Grammer, R., Gross, W.L., Urman, R.D. **Analysis of adverse events associated with adult moderate procedural sedation outside the operating room.** *J Patient Saf.* 2014; ([Epub ahead of print]).

[Google Scholar](#)

18. Murphy, L.A., Thornburg, L.L., Glantz, J.C., Wasserman, E.C., Stanwood, N.L., Betstadt, S.J.

Complications of surgical termination of second-trimester pregnancy in obese versus nonobese women. *Contraception.* 2012;86:402–406.

[Abstract](#) | [Full Text](#) | [Full Text PDF](#) | [PubMed](#) | [Scopus \(7\)](#) | [Google Scholar](#)

19. Gokhale, P., Lappen, J.R., Waters, J.H., Perriera, L.K. **Intravenous sedation without intubation and the risk of anesthesia complications for obese and non-obese women undergoing surgical abortion: a retrospective cohort study.** *Anesth Analg.* 2016;122:1957–1962.

[Crossref](#) | [PubMed](#) | [Scopus \(1\)](#) | [Google Scholar](#)

20. Bartlett, L.A., Berg, C.J., Shulman, H.B., Zane, S.B., Green, C.A., Whitehead, S. et al, **Risk factors for legal induced abortion-related mortality in the United States.** *Obstet Gynecol.* 2004;103:729–737.

[Crossref](#) | [PubMed](#) | [Scopus \(232\)](#) | [Google Scholar](#)

21. Rausch, M., Lorch, S., Chung, K., Frederick, M., Zhang, J., Barnhart, K. **A cost-effectiveness analysis of surgical versus medical management of early pregnancy loss.** *Fertil Steril.* 2012;97:355–360.

[Abstract](#) | [Full Text](#) | [Full Text PDF](#) | [PubMed](#) | [Scopus \(28\)](#) | [Google Scholar](#)

☆Sponsor: Society of Family Planning Research Fund (SFPRF).

☆Harvard Medical School Center for Primary Care

☆Harvard Medical School Scholars in Medicine Program

☆☆Disclosures: The authors have no conflicts of interest. The findings and conclusions expressed in this article are those of the authors and do not necessarily reflect the views of Planned Parenthood Federation of America, Inc.

★Presented at: Oral Presentation, North American Forum on Family Planning, Denver, CO, Nov. 5, 2016.

© 2017 Elsevier Inc. All rights reserved.

[< Previous Article](#)

January 2018 Volume 97, Issue 1, Pages 48–53

[Next Article >](#)

