

Laminaria Insertion & Induction of Intrauterine Fetal Demise.

Patient's Name: [REDACTED] Date: 8/12/10 Chart#: 1562

Vital Signs: BP: 104/70 Pulse: 85 Temp: 97.4

I have discussed with the patient the abortion she has requested and I believe she is sufficiently mature and intelligent to understand the nature and consequences of her decision.

The patient was placed in the lithotomy position. The perineum was prepped and draped.

On Pelvic Examination:

Vagina [X] WNL [] Other: _____
Cervix [X] WNL [] Other: _____
Adnexa [X] WNL [] Other: _____
Uterus [X] WNL [] Other: _____ [] Ant [] Mid [] Post

Approximate Size of Uterus:

13 14 15 16 17 18 19 20 21 22 23 24 25

[] Periumbilical prep was done. 10cc of 1% Lidocaine injected subcutaneously. Under ultrasonographic guidance a 7-inch spinal needle was inserted through the skin and into the fetus.

_____ cc. (500mcg/2cc) Digoxin was injected to cause fetal demise.

Laminaria insertion: Total number of Laminaria Inserted: 7 LOCAL/TWILIGHT

- 5 10mm Laminaria Japonica [] Membranes ruptured.
- _____ 8mm Laminaria Japonica [] Umbilical chord ligated.
- _____ 6mm Laminaria Japonica [] Lamicol 5mm. # _____
- _____ 5mm Laminaria Japonica [] Lamicol 3mm. # _____
- _____ 4mm Laminaria Japonica
- _____ 3mm Laminaria Japonica
- 2 2mm Laminaria Japonica

Sterile gauze packing was then placed in the vagina. Patient was taken to the recovery room for observation.

Medications administered and/or prescribed to her:

- _____ Misoprostol tabs, _____ x 200 mg, p.o. started at _____ am/pm. and repeated every _____ hrs.
- _____ Misoprostol tabs (_____ #) placed p.v.
- _____ Mifepristone tabs 100 mg p.o. at _____
- _____ Hydrocodone
- _____ Dilaudid
- X Doxycycline 100 mg B.I.D.
- X Other: Tylenol # 3

* <

Return to Vauh office on 8/13 at 8 am/pm.

MD signature: [Signature]