

Exhibit A

Abortion Procedure Record
VII-PPHP-2

Staple Sono Pictures to Back of Form

Staple Laminaria Wrappers to Back of Form

Planned Parenthood®
Hudson Peconic, Inc.
Serving Suffolk, Westchester, Rockland and Putnam Counties

DEJESUS, SHARA
Acct#: 889905
DOB: 01/15/1967

ALLERGIES:
List: None
 NKDA

ABORTION PROCEDURE RECORD

Con.Sed.
Local
NPO ≥ 6 hrs
Yes No

LMP: 11/13/10 G. A. by Sono 6-4 Date of Procedure: 2/27/16
Age: 43 Preg: 5 #Deliv: 2 #SuAb: 1 #MedAB: 5 #Miscarriage: 1 #Ectopic: 2 #Living Children: 2
Rh: + Hct/Hbg: 10.4 Temp: 99.2 BP: 135/85 Pulse: 68

Height: 5'7" Weight: 246 BCM Desired: Condoms
C-Section history: Yes No number of c-sections: 1
Asthma: Yes No
Breastfeeding: Yes No
Pertinent medical and surgical history: Q

Yes No
 AB consent signed
 Birth Control info given
 Under 18 counseling

No Pre-op Sodium 500 mg po - given @ _____ by _____
Lab Staff Signature: [Signature] Counselor Signature: [Signature]

CERVICAL PREPARATION:

Consents signed for cervical preparation and surgical procedure
 Medical History reviewed/patient cleared for surgery
Laminaria:
 Naproxen 500 mg po - given @ _____ by _____
The cervix was cleansed and _____ cc 1% lidocaine was instilled for paracervical block. A tenaculum was applied and _____ No. laminaria and _____ sponges were inserted. Patient tolerated procedure well.
Comments:
 Post instructions given. Doxycycline 100mg PO BID x 7 days (start now)
Laminaria inserted by _____ Date/Time: _____
Misoprostol:
_____ mcg misoprostol was given buccally at _____ AM/PM
Misoprostol given by: _____ Date: _____

Sonogram: Ultrasound and report reviewed
Consents: I have reviewed the consent forms and answered all patient questions.
Exam: Abdomen: nl Vulva: nl Vag: nl Cervix: nl Adnexa: nl
GC/CHL done Uterus: 0 weeks, position: Anterior
Intent: I intend to utilize disarticulation/dismemberment techniques to remove the fetus using suction and/or forceps.
Procedure: The cervix was cleansed and tenaculum applied. _____ laminaria and _____ sponges removed. 14 cc of 1% lidocaine (with _____ ju vasopressin) was instilled as a paracervical block. The cervix was dilated to 14 Pratt and 6 mm cannula was used to empty the uterus using Electromechanical suction or MVA. Sharp curettage was performed.

The POC/fetus was removed using forceps in addition to suction. Fetal demise occurred prior to the fetus reaching the following landmarks: if head first, before the entire fetal head was outside the body of the woman, and if breech, before any part of the fetal trunk past the navel was outside the body of the woman.
 multiple passes of forceps occurred resulting in extraction of the fetus in multiple parts
 the umbilical cord was severed before removal of fetal parts
 the umbilical cord was not pulsating prior to evacuation to the anatomical landmarks listed above.

Sono guidance was used during procedure. Sono post-operatively demonstrated that the uterus was empty.
Uterotonics administered: Methergine 0.2 mg IM x 1 Methergine 0.2 mg IC x 1 Oxytocin _____ mAU per Liter
 The patient was transferred to the recovery room in stable condition.
Comments:
Estimated blood loss: 210 cc
Physician Signature: [Signature]