

We use cookies to enhance your experience on our website. By clicking 'continue' or by continuing to use our website, you are agreeing to our use of cookies. You can change your cookie settings at any time.

[Continue](#)[Find out more](#)

Birth outcomes after induced abortion: a nationwide register-based study of first births in Finland FREE

R. Klemetti , M. Gissler, M. Niinimäki, E. Hemminki

Human Reproduction, Volume 27, Issue 11, November 2012, Pages 3315–3320,

<https://doi.org/10.1093/humrep/des294>

Published: 29 August 2012 **Article history** ▼

[PDF](#)[Split View](#)[Cite](#)[Permissions](#)[Share](#) ▼

Abstract

STUDY QUESTION

Is the perinatal health of first-born children affected by the mother's previous induced abortion(s) (IAs)?

SUMMARY ANSWER

Prior IAs, particularly repeat IAs, are correlated with an increased risk of some health problems at first birth; even in a country with good health care quality.

WHAT IS KNOWN ALREADY

A positive association between IA and risk of preterm birth or a dose–response effect has been found in some previous studies. Limited information and conflicting results on other infant outcomes are available.

Sk

STUDY DESIGN, SIZE AND DURATION

[Article Navigation](#)

PARTICIPANTS/MATERIALS, SETTING AND METHODS

All the first-time mothers with a singleton birth (obtained from the Medical Birth Register) in the period 1996–2008 ($n = 300\,858$) were linked to the Abortion Register for the period 1983–2008.

MAIN RESULTS AND THE ROLE OF CHANCE

Of the first-time mothers, 10.3% ($n = 31\,083$) had one, 1.5% had two and 0.3% had three or more IAs. Most IAs were surgical (88%) performed before 12 weeks (91%) and carried out for social reasons (97%). After adjustment, perinatal deaths and very preterm birth (<28 gestational week) suggested worse outcomes after IA. Increased odds for very preterm birth were seen in all the subgroups and exhibited a dose–response relationship: 1.19 [95% confidence interval (CI) 0.98–1.44] after one IA, 1.69 (1.14–2.51) after two and 2.78 (1.48–5.24) after three IAs. Increased odds for preterm birth (<37 weeks) and low birthweight (<2500 g and <1500 g) were seen only among mothers with three or more IAs: 1.35 (1.07–1.71), 1.43 (1.12–1.84) and 2.25 (1.43–3.52), respectively.

LIMITATIONS, REASONS FOR CAUTION

Observational studies like ours, however large and well-controlled, will not prove causality.

WIDER IMPLICATIONS OF THE FINDINGS

In terms of public health and practical implications, health education should contain information of the potential health hazards of repeat IAs, including very preterm birth and low birthweight in subsequent pregnancies. Health care professionals should be informed about the potential risks of repeat IAs on infant outcomes in subsequent pregnancy.

STUDY FUNDING/COMPETING INTEREST(S)

National Institute for Health and Welfare and the Academy of Finland. No competing interests.

Keywords: [induced abortion](#), [termination of pregnancy](#), [birth outcomes](#), [prematurity](#), [low birthweight](#), [register study](#)

Topic: [pregnancy](#), [quality of care](#), [conflict of interest](#), [abortion](#), [induced](#), [child](#), [finland](#), [health personnel](#), [infant](#), [low birth weight infant](#), [mothers](#), [personal satisfaction](#), [social role](#),

Issue Section: [Reproductive Epidemiology](#)

Introduction

Induced abortion (IA) is one of the most common gynecological procedures performed on women but the impact of IAs on women's subsequent childbearing has not yet been properly studied ([van Oppenraaij *et al.*, 2009](#); [Lowit *et al.*, 2010](#)).

Many previous studies ([Henriet and Kaminski, 2001](#); [Ancel *et al.*, 2004](#); [Ngyen *et al.*, 2004](#); [Moreau *et al.*, 2005](#); [Freak-Poli *et al.*, 2009](#); [Voigt *et al.*, 2009](#)) and reviews ([van Oppenraaij *et al.*, 2009](#); [Shah and Zao, 2009](#); [Swingle *et al.*, 2009](#); [Lowit *et al.*, 2010](#)) have found a positive association between IA and risk of preterm birth or a dose–response effect (i.e. the risk of preterm birth increased with increasing number of IAs). Some smaller studies did not find the association ([Reime *et al.*, 2008](#)) or it disappeared after adjustment for confounding factors ([Raatikainen *et al.*, 2006](#)). The limited data on the association between low birthweight and pre-birth IA have produced conflicting results ([Zhou *et al.*, 2000](#); [Henriet and Kaminski, 2001](#); [Raatikainen *et al.*, 2006](#); [Parazzini *et al.*, 2007](#); [Reime *et al.*, 2008](#)).

Most of the recent original studies have been based on small sample-sizes ([Henriet and Kaminski, 2001](#); [Ancel *et al.*, 2004](#); [Ngyen *et al.*, 2004](#); [Raatikainen *et al.*, 2006](#); [Parazzini *et al.*, 2007](#); [Reime *et al.*, 2008](#)) and on women's self-reports on exposure ([Henriet and Kaminski, 2001](#); [Ancel *et al.*, 2004](#); [Ngyen *et al.*, 2004](#); [Moreau *et al.*, 2005](#); [Raatikainen *et al.*, 2006](#); [Parazzini *et al.*, 2007](#); [Reime *et al.*, 2008](#); [Freak-Poli *et al.*, 2009](#); [Voigt *et al.*, 2009](#)), while the most important confounding factors have not always been controlled for ([Zhou *et al.*, 2000](#); [Henriet and Kaminski, 2001](#); [Ancel *et al.*, 2004](#); [Moreau *et al.*, 2005](#); [Raatikainen *et al.*, 2006](#); [Parazzini *et al.*, 2007](#); [Reime *et al.*, 2008](#); [Freak-Poli *et al.*, 2009](#); [Voigt *et al.*, 2009](#)). Furthermore, the type of procedures used, the gestational age at the time of abortion and the number of previous IAs vary substantially ([Atrash and Hogue, 1990](#)). Thus, there is a clear need for further studies, particularly from countries with high-quality IA services and reliable information on IAs. Compared with other European countries, Finland has a low rate of IA: 8.9 per 1000 women aged 15–49 ([Gissler *et al.*, 2011](#)). According to Finnish legislation enacted in 1970 with updates in 1978 and 1985, a woman needs permission with a legal indication (social, medical or ethical reasons) for IA, although the legislation is interpreted liberally up to 12 weeks. An IA can be legally performed up to 20 weeks, and in cases of a

[Article Navigation](#)

The purpose of this study was to examine birth outcomes of first-born children by mother's IA history in Finland, which has good IA and birth services and relatively few IAs. Our hypothesis was that having one IA does not affect birth outcomes, but having several may result in poorer birth outcomes. The study was based on nationwide obligatory health registers and took into account mothers' background characteristics, previous miscarriages and ectopic pregnancies.

Materials and Methods

All first-time mothers with a singleton birth were identified in the nationwide Finnish Medical Birth Register (MBR) for the period 1996–2008 and linked to the data in the Abortion Register (AR) for the period 1983–2008. The MBR was started in 1987 and includes all births occurring in Finland (THL, 2012a). The register contains information on mothers' background, care during pregnancy and delivery and infants' health up to the age of 7 days. The AR has been operating since 1950, and computerized data have been available since 1983 (Gissler *et al.*, 1996, 2004). The AR is based on obligatory notification from physicians (THL, 2012b). The AR includes information on reasons for IA, date, method and gestational weeks of IA, as well as information on woman's sociodemographic and obstetric background. According to data quality studies, the coverage and accuracy of both registers are very good and the contents correspond well to the information found in medical records (Gissler *et al.*, 1995, 1996).

Mother's first birth was identified from the MBR, and her IAs prior to that date were checked from the AR (notifications) and the MBR (mothers' own information). Mothers were classified according to the information in the AR by the number of IAs before the first birth: no IA, one IA, two IAs and three or more IAs. Mothers who, according to the MBR, had an IA history but in whom no IAs were found in the AR were classified as 'IA history unknown'. If there was a discrepancy in the number of IAs between the MBR and AR, the AR information was used ($n = 5950$).

Birth outcomes by the mother's number of IAs were calculated. The following were used as outcome indicators: very low birthweight (<1500 g), low birthweight (<2500 g), very preterm birth (<28 weeks), preterm birth (<37 weeks), low one-minute Apgar scores (0–6) and perinatal death (from 22 weeks). Differences in mothers' background characteristics were studied by cross tabulations and chi-square tests. In the MBR, the socioeconomic position of the mother is defined by using her own occupation at the time of delivery, which is collected

[Skip to Main Content](#)

Article Navigation

classification into four categories was used: upper white-collar workers, lower white-collar workers, blue-collar workers and others (entrepreneurs, students, pensioners, unemployed women and women with an unclassified position).

Logistic regression (odds ratios and 95% confidence intervals) was used to adjust for background variables considered confounders. Confounders were selected on the basis of previous literature on maternal risk factors on birth outcomes and their availability and quality in the used health registers (Gissler *et al.*, 1995, 1996). The original confounder list included maternal age, marital status, socioeconomic position, urbanity, smoking during pregnancy, previous ectopic pregnancies and miscarriages, as well as method, indication and timing of IA. Of those, only statistically significant variables ($P < 0.1$) were included in the model. All the variables related to IAs (method, indication and timing) were excluded, since their inclusion did not improve the model. First, adjustment was made for social background (maternal age, marital status, socioeconomic position, urbanity and smoking during pregnancy). Second, data were adjusted for previous ectopic pregnancies and miscarriages. Mothers with an unknown history of IAs were excluded from the logistic regression analyses. Statistical analyses were performed by SAS, version 9.1.

Ethical approval for the study was obtained from the Ethics committee of the National Institute for Health and Welfare (THL), while THL also gave permission to use the registers in this study.

Results

In total, 300 858 first-time mothers with singleton births in the period 1996–2008 were identified from the MBR. According to the AR, 31 083 (10.3%) had had one, 4417 (1.5%) two and 942 (0.3%) three or more IAs before the first birth; 226 mothers (0.08%) had an unknown history of IA. Of IAs, 88% were surgical, 91% were made before 12 weeks and 97% were made for social reasons.

Compared with mothers with no previous IAs, mothers with previous IAs were more often smokers, single, from urban areas and from a lower socioeconomic position, and had had miscarriages and ectopic pregnancies before their first birth (Table I).

[Skip to Main Content](#)

Table I

[Article Navigation](#)
Background variable^b**Number of previous induced abortions**

	0 (n = 264 190)		1 (n = 31 083)		2 (n = 4417)		3+ (n = 942)		Unknown history^c (n = 226)		Total (n = 300 858)	
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
Age group												
< 20 years	15 759	6.0	2411	7.8	249	5.6	26	2.8	10	4.4	18 455	6.1
20–24	66 434	25.1	8668	27.9	1128	25.5	194	20.6	48	21.2	76 472	25.4
25–29	98 392	37.2	9613	30.9	1320	29.9	295	31.3	77	34.1	109 697	36.5
30–34	60 284	22.8	6805	21.9	1069	24.2	249	26.4	59	26.1	68 466	22.7
35–39	19 601	7.4	2949	9.5	534	12.1	133	14.1	25	11.1	23 242	7.7
40+	3719	1.4	637	2.0	117	2.6	45	4.8	7	3.1	4525	1.5
Unknown	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.0
Area												
Urban	1 84 926	70.0	22 591	72.7	3365	76.2	720	76.4	109	48.2	2 11 711	70.4
Semi-urban	39 389	14.9	4251	13.7	517	11.7	104	11.0	70	31.0	44 331	14.7
Rural	39 405	14.9	4209	13.5	519	11.8	116	12.3	36	15.9	44 285	14.7
Abroad	470	0.2	32	0.1	16	0.4	2	0.2	11	4.9	531	0.2
Marital status												

[Skip to Main Content](#)

Article Navigation

	0 (n= 264 190)		1 (n= 31 083)		2 (n= 4417)		3+ (n= 942)		Unknown history^c (n = 226)		Total (n = 300 858)	
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
Married	1 29 295	48.9	9785	31.5	1188	26.9	272	28.9	86	38.1	1 40 626	46.7
Cohabiting	93 223	35.3	13 813	44.4	1972	44.6	365	38.7	112	49.6	1 09 485	36.4
Single	32 654	12.4	6202	20.0	1092	24.7	262	27.8	22	9.7	40 232	13.4
Unknown	9018	3.4	1283	4.1	165	3.7	43	4.6	6	2.7	10 515	3.5
Socioeconomic position												
Upper white-collar	42 551	16.1	3211	10.3	362	8.2	73	7.7	21	9.3	46 218	18.3
Lower white-collar	79 354	30.0	9101	29.3	1223	27.7	245	26.0	47	20.8	89 970	35.6
Blue-collar	30 749	11.6	5169	16.6	840	19.0	178	18.9	20	8.8	36 956	14.6
Other ^d	1 11 536	42.2	13 602	43.8	1992	45.1	446	47.3	138	61.1	1 27 714	31.4
Smoker ^e	39 200	14.8	9285	29.9	1821	41.2	433	46.0	48	21.2	50 787	16.9
History of miscarriage	26 476	10.0	3671	11.8	613	13.9	129	13.7	21	9.3	30 910	10.3
History of ectopic pregnancy	2533	1.0	418	1.3	87	2.0	21	2.2	1	0.4	3060	1.0

[Skip to Main Content](#)

^aInduced abortions in the AR during 1983–2008.

[Article Navigation](#)

^cHistory of induced abortion in the Medical Birth Register but no induced abortions in the AR.

^dOther = entrepreneurs, students, pensioners, unemployed women and women with an unclassified position.

^eSmoking during pregnancy.

[Open in new tab](#)

In the unadjusted analysis, all poor birth outcomes increased by the number of IAs (Table II). Poor outcomes were rare among the mothers with an unknown history of IAs, with numbers being too low to judge their risk.

Table II

Birth outcomes of first-time Finnish mothers during 1996–2008 by the number of previous induced abortions^a.

Outcome	Number of previous induced abortions										
	0 (n = 264 190)		1 (n = 31 083)		2 (n = 4417)		3+ (n = 942)		Unknown history ^b (n = 226)		Total (n = 858)
	n	/1000 ^c	n	/1000 ^c	n	/1000 ^c	n	/1000 ^c	n	/1000 ^c	n
Very preterm <28 gestational week	837	3	120	4	26	6	10	11	0	0	993
Preterm <37 gestational week	14 705	56	1737	60	269	63	77	84	8	35	16 796
Very low birth weight <1500 g	2087	8	266	9	46	11	20	22	0	0	2419

[Skip to Main Content](#)

Article Navigation

	0 (<i>n</i> = 264 190)		1 (<i>n</i> = 31 083)		2 (<i>n</i> = 4417)		3+ (<i>n</i> = 942)		Unknown history ^b (<i>n</i> = 226)		Total 858)
	<i>n</i>	/1000 ^c	<i>n</i>	/1000 ^c	<i>n</i>	/1000 ^c	<i>n</i>	/1000 ^c	<i>n</i>	/1000 ^c	<i>n</i>
Low birthweight <2500 g	11 097	42	1353	47	224	52	70	76	8	35	12 752
Apgar scores 1 min 0–6	17 595	67	1997	69	317	74	73	80	17	75	19 999
Perinatal death	1272	5	187	6	29	7	9	10	1	4	1498

^aInduced abortions in the AR during 1983–2008.

^bHistory of induced abortion in the Medical Birth Register but no induced abortions in the AR.

^cNumber per 1000 births.

[Open in new tab](#)

After adjustment for social background, very preterm birth (<28 gestational week) suggested worse outcomes (Table III). By the number of IAs, the risk of very preterm birth was seen in all the subgroups (even though not always statistically significant), as was a dose–response effect (Table III). Risk of very low birthweight (<1500 g) was similar, but was statistically significant only among women with three or more IAs. Risks of preterm birth (<37 weeks) and low birthweight (<2500 g) were seen only among mothers with three or more IAs. Low Apgar scores were not related to IAs. Results on perinatal deaths suggested a small increase after IA, but most comparisons were statistically non-significant (Table III). The largest increase was after three IAs.

[Skip to Main Content](#)

Table III

[Article Navigation](#)

abortions, by the number of abortions.

Birth outcome	Crude and adjusted odds ratios and 95% CI							
	0 versus 1		0 versus 2		0 versus 3+		0 versus 1+	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
Very preterm <28 gestational week								
Crude Model	1.22	1.01–1.48	1.86	1.26–2.76	3.38	1.80–6.32	1.34	1.13–1.60
Adjusted Model I ^a	1.20	0.99–1.45	1.71	1.15–2.54	2.81	1.49–5.29	1.28	1.07–1.53
Adjusted Model II ^b	1.19	0.98–1.44	1.69	1.14–2.51	2.78	1.48–5.24	1.27	1.06–1.52
Preterm <37 gestational week								
Crude Model	1.00	0.95–1.06	1.10	0.97–1.25	1.51	1.20–1.91	1.03	0.98–1.08
Adjusted Model I ^a	0.98	0.93–1.03	1.02	0.90–1.15	1.36	1.07–1.72	0.99	0.95–1.04
Adjusted Model II ^b	0.98	0.93–1.03	1.01	0.89–1.15	1.35	1.07–1.71	0.99	0.94–1.04
Very low birthweight <1500 g								
Crude Model	1.08	0.95–1.23	1.32	0.99–1.77	2.72	1.75–4.25	1.15	1.02–1.29
Adjusted Model I ^a	1.04	0.91–1.18	1.14	0.84–1.54	2.26	1.44–3.53	1.06	0.94–1.20
Adjusted Model II ^b	1.03	0.91–1.18	1.13	0.84–1.54	2.25	1.43–3.52	1.06	0.94–1.19
Low birthweight <2500 g								
Crude Model	1.04	0.98–1.10	1.22	1.06–1.40	1.83	1.43–2.34	1.08	1.02–1.14
Adjusted Model I ^a	0.96	0.90–1.01	1.02	0.89–1.18	1.44	1.12–1.84	0.98	0.93–1.03
Adjusted Model II ^b	0.96	0.90–1.01	1.02	0.89–1.18	1.43	1.12–1.84	0.98	0.93–1.03
Apgar scores 1 min 0–6								
Crude Model	0.96	0.92–1.01	1.08	0.97–1.22	1.18	0.93–1.50	0.98	0.94–1.03

[Skip to Main Content](#)

[Article Navigation](#)

	0 versus 1		0 versus 2		0 versus 3+		0 versus 1+	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
Adjusted Model I ^a	0.95	0.90–1.00	1.05	0.93–1.18	1.11	0.87–1.41	0.97	0.93–1.02
Adjusted Model II ^b	0.95	0.90–1.00	1.05	0.93–1.18	1.10	0.86–1.41	0.97	0.93–1.02
Perinatal death								
Crude Model	1.25	1.07–1.46	1.37	0.94–1.98	1.99	1.03–3.85	1.28	1.11–1.48
Adjusted Model I ^a	1.19	1.02–1.39	1.16	0.79–1.71	1.70	0.88–3.30	1.15	0.99–1.34
Adjusted Model I ^c	1.15	0.98–1.36	1.16	0.79–1.71	1.70	0.87–3.29	1.17	1.01–1.36
Adjusted Model II ^b	1.19	1.02–1.39	1.16	0.79–1.70	1.70	0.87–3.29	1.15	0.99–1.34

^aAdjusted for social factors: age, marital status, socioeconomic position, urbanity, smoking; women with missing information in the AR ($n = 226$, 0.08%) excluded.

^bAdjusted additionally for miscarriage and ectopic pregnancy; women with missing information in the AR ($n = 226$, 0.08%) excluded.

^cExcluding women who had a history of induced abortion in the Medical Birth Register but not in the AR ($n = 226$) and those with a discrepancy in the number of induced abortions between the Medical Birth Register and the AR ($n = 5950$).

No induced abortion, $n = 264\,190$ (a reference group), 1 induced abortion $n = 31\,083$, 2 induced abortions $n = 4417$, 3+ induced abortions $n = 942$, 1 or more induced abortions, $n = 36\,442$.

[Open in new tab](#)

Because the overall risk of perinatal death was lower than risks in each subgroup, we also did a sub-analysis excluding the women who had a discrepancy in the number of abortions recorded in the AR and the MBR (Table III). The results remained similar.

[Skip to Main Content](#)

Further adjustment for history of miscarriage and ectopic pregnancy had only a marginal effect on adverse birth outcomes after IAs (Table III).

[Article Navigation](#)

In our study, after adjusting for mothers' background characteristics, mothers who had had one pre-birth IA had only marginally increased risks of some outcomes. After two previous IAs, the risk of very preterm birth was increased statistically significantly. With the exception of low Apgar scores, the studied adverse birth outcomes were more common among mothers having had three or more IAs. Taking into account previous miscarriages and ectopic pregnancies did not notably change the results.

Owing to the large number of IAs carried out every year, even a very small increase in risk of poor birth outcomes could have significant public health implications. The majority of previous studies of birth outcomes after pre-birth IA have had methodological problems. Prior to our study, only one large-scale study with reliable data on the exposure (IA) in a country with good IA and birth services and relatively few IAs (Denmark) had been published ([Zhou *et al.*, 2000](#)). Our study was the first that can at least partly control for the most important confounding factors, such as smoking and socioeconomic position. Furthermore, we were able to study more outcomes than just the low-birthweight variable investigated in the Danish study ([Zhou *et al.*, 2000](#)).

Our large national register-based study covered all the first-time mothers having a singleton birth in the period 1996–2008 and all the IAs carried out in Finland in the period 1983–2008. Data on birth outcomes received from the MBR are considered reliable ([Gissler *et al.*, 1995](#)). We did not have information on IA before 1982 or on IAs performed in other countries. Earlier studies assessing the completeness of the Finnish AR found that 99% of IAs were reported to the register and at least 95% of the information matched with medical records ([Gissler *et al.*, 1996](#); [Heikinheimo *et al.*, 2008](#)). We found 226 first-time mothers—0.08% of all the studied first-time mothers—in the MBR whose IA history was unknown. As the IA information in the MBR is originally based on mothers' information, some of these may be miscarriages; others may have been mistakes or IAs done while the woman was outside Finland or prior to 1983. Poor outcomes were rare among these mothers. Women who had a discrepancy in the number of IAs between the registers were classified according to the AR and analyses were made that both included and excluded them, which had only marginal effects on the results.

As in previous studies ([Ratikainen *et al.*, 2006](#); [Heikinheimo *et al.*, 2008](#); [Niinimäki *et al.*, 2009](#)), we found that the demographic and reproductive profiles of first-time mothers who had had a pre-birth IA differed from those of the first-time mothers who had had no pre-birth IAs. Most risk factors such as smoking, lower socioeconomic position and previous miscarriages and ectopic pregnancies were more common among primiparous women who had experienced IA. In the logistic regression analyses, we were able to adjust for these

[Skip to Main Content](#)

Article Navigation

with low socioeconomic position, which is also a risk factor for prematurity and perinatal mortality. Most IAs in our study were made during the first trimester (91%).

In contrast to some earlier studies ([Ancel et al., 2004](#); [Moreau et al., 2005](#); [Freak-Poli et al., 2009](#); [Voigt et al., 2009](#)), we did not find any significant increased risk of preterm birth after one IA ([Henriet and Kaminski, 2001](#); [Ancel et al., 2004](#); [Freak-Poli et al., 2009](#); [Shah and Zao, 2009](#); [Voigt et al., 2009](#)). Neither did we find a dose–response effect between the number of pre–birth IAs and preterm birth. An increased risk of preterm birth was found only after three or more pre–birth IAs. In keeping with some earlier findings, we found a dose–response effect between the number of pre–birth IAs and very preterm birth ([Moreau et al., 2005](#); [Freak-Poli et al., 2009](#); [Shah and Zao, 2009](#); [Voigt et al., 2009](#)).

Our study's finding of a stronger association of pre–birth IA with very preterm birth than with preterm birth is in line with the findings of some earlier studies ([Lumley, 1998](#); [Ancel et al., 2004](#); [Moreau et al., 2005](#)). It has been suggested that both infections before and after IA and surgical procedures may be the underlying mechanisms for the increased risk of very preterm births in subsequent pregnancies ([Henriet and Kaminski, 2001](#); [Ancel et al., 2004](#)). Women with a history of IA have had an increased risk of intra–amniotic infection, intra–partum infection and infections of their infants. A relationship has been found between previous IA and preterm birth after placenta previa and other maternal hemorrhage ([Ancel et al., 2004](#)), and it has been suggested that the surgical IA, by damaging the endometrium, may increase faulty placentation, causing preterm delivery in subsequent pregnancy. Furthermore, surgical IA may cause mechanical trauma to the cervix, increasing the risk of cervical insufficiency ([Ancel et al., 2004](#)). In repeat IAs, women are repeatedly exposed to these potentially harmful effects.

Eighty–eight percent of the IAs in this study were done surgically. The women with three or more IAs had twice as many (2.2 versus 1.0%) ectopic pregnancies as there were in women with no IAs. This may reflect a previous history of pelvic inflammatory disease and Chlamydia infections, which are also risk factors for preterm birth ([Paavonen, 2012](#)). However, adjusting for previous miscarriages or ectopic pregnancies did not notably change our findings.

Preterm birth has been the most studied outcome, and there are very few data on other birth outcomes ([van Oppenraaij et al., 2009](#)). As with the Danish study ([Zhou et al., 2000](#)), we could not find a strong association between low birthweight and a previous IA, although in contrast to the Danish study, we did find an increased risk after three or more previous IAs and a dose–response effect in very low birthweight as described in one previous meta–analysis

Article Navigation

(van Oppenraaij *et al.*, 2009). We found the highest risk after three or more IAs, but even our sample was not large enough to enable statistical significance. Our result of a non-existing association between low Apgar scores and pre-birth IA(s) is in line with the findings of previous studies (van Oppenraaij *et al.*, 2009).

Only a few studies have taken into account the method of IA when studying the association between poor birth outcomes and pre-birth IAs, and those taking it into account did not find any differences by the IA method (Atrash and Hogue, 1990; Yimin *et al.*, 2003; Virk *et al.*, 2007). According to a recent Finnish study, medical IA offered a good alternative to surgical methods without increasing the risk of repeat IAs, but with an increased risk of short-term adverse events (Niinimäki *et al.*, 2009). In 2010, the majority (90%) of IAs in Finland were carried out by medical methods (THL, 2012b). During our study period (1983–2008) most IAs were carried out by surgical methods. In the future, when the number of IAs carried out by medical methods is high enough (but surgical IAs still exist), a large-scale study should be done that compares the birth outcomes in the subsequent pregnancy by the method of IA.

In conclusion, our study confirms those previous findings suggesting that prior repeat IAs are correlated with an increased the risk of (very) preterm birth and (very) low birthweight at first birth. From a public health perspective, information on increased risk of very preterm birth after IAs and its dose–response effect is intrinsically important. In terms of practical implications, it must be noted that observational studies like ours, however large and well-controlled, will not prove causality. For example, our finding of an increased risk following several IAs may be due to some confounding that we could not control for. However, as there are also other reasons to avoid IAs and particularly repeat IAs, sexuality education should contain information of the potential health hazards of IAs, including for subsequent pregnancies. Health care professionals should be informed about the potential risks of repeat IAs on birth outcomes in subsequent pregnancy.

Authors' roles

R.K.: participated in the interpretation of data, drafting and design of the article and in the modification of the final version. M.G.: participated in the interpretation and analysis of data, planning the article, revising the draft critically and approving the final version. M.N.: participated in the interpretation of data, revising the draft critically and approving the final version. E.H.: participated in the conception and design of the article, interpretation of data, revising the draft critically and approving the final version.

[Article Navigation](#)

This study was supported by the National Institute for Health and Welfare and the Academy of Finland.

Conflict of interest

None declared.

References

Ancel PY, Lelong N, Papiernik E, Áphe Saurel-Cubizolles MJ, Kaminski M. History of induced abortion as a risk factor for preterm birth in European countries: results of the EUROPOP survey, *Hum Reprod*, 2004, vol. 19 (pg. 734-740)[10.1093/humrep/deh107](#).

[Google Scholar](#) [Crossref](#) [PubMed](#). [Crossref](#)

Atrash HK, Hogue CJ. The effect of pregnancy termination on future reproduction, *Baillieres Clin Obstet Gynecol*, 1990, vol. 4 (pg. 391-405)

[Google Scholar](#) [Crossref](#)

Freak-Poli R, Chan A, Tucker G, Street J. Previous abortion and risk of pre-term birth: a population study, *J Matern Fetal Neonatal Med*, 2009, vol. 22 (pg. 1-7)

[Google Scholar](#) [Crossref](#) [PubMed](#).

Gissler M, Teperi J, Hemminki E, Meriläinen J. Data quality after restructuring a nationwide medical birth registry, *Scand J Soc Med*, 1995, vol. 23 (pg. 75-80)

[Google Scholar](#) [PubMed](#).

Gissler M, Ulander VM, Hemminki E, Rasimus A. Declining induced abortion rate in Finland: data quality of the Finnish abortion register, *Int J Epidemiol*, 1996, vol. 25 (pg. 376-380)

[Google Scholar](#) [Crossref](#) [PubMed](#).

[Skip to Main Content](#)

Gissler M, Berg C, Bouvier-Colle MH, Buekens P. Pregnancy-associated mortality after birth, spontaneous abortion, or induced abortion in Finland, 1987–2000, *Am J Obstet*

[Article Navigation](#)

Gissler M, Fronteira I, Jahn A, Karro H, Moreau C, Oliveira da Silva M, Olsen J, Savona-Ventura C, Temmerman M, Hemminki E. the REPROSTAT group Terminations of pregnancy in the European Union, *Br J Obstet Gynaecol*, 2011 . (Corrigendum: doi: 10.1111/j.1471-0528.2012.03290.x)[10.1111/j.1471-0528.2011.03189.x](https://doi.org/10.1111/j.1471-0528.2011.03189.x).

[Crossref](#)

Heikinheimo O, Gissler M, Suhonen S. Age, parity, history of abortion and contraceptive choices affect the risk of repeat abortion, *Contraception*, 2008, vol. 78 (pg. 149-154)

[Google Scholar](#) [Crossref](#) [PubMed](#).

Henriet L, Kaminski M. Impact of induced abortion on subsequent pregnancy outcome: the 1995 French national perinatal survey, *Br J Obstet Gynaecol*, 2001, vol. 108 (pg. 1036-1042)

Lowit A, Bhattacharya S, Bhattacharya S. Obstetric performance following an induced abortion, *Best Pract Res Clin Obstet Gynaecol*, 2010, vol. 24 (pg. 667-682)

[Google Scholar](#) [Crossref](#) [PubMed](#).

Lumley J. The association between prior spontaneous abortion, prior induced abortion and preterm birth in first singleton births, *Prenat Neonat Med*, 1998, vol. 3 (pg. 21-24)

Moreau C, Kaminski M, Ancel PY, Bouyer J, Escande B, Thiriez G, Boulot P, Fresson J, Arnaud C, Subtil D, et al. EPIPAGE Group Previous induced abortions and the risk of very preterm delivery: results of the EPIPAGE study, *Br J Obstet Gynaecol*, 2005, vol. 112 (pg. 430-437)

[Google Scholar](#) [Crossref](#)

Ngyen N, Savitz DA, Thorp JM. Risk factors for preterm birth in Vietnam, *Int J Gynecol Obstet*, 2004, vol. 86 (pg. 70-78)

[Google Scholar](#) [Crossref](#)

[Skip to Main Content](#)
Niinimäki M, Pouta A, Bloigu A, Gissler M, Hemminki E, Suhonen S, Heikinheimo O. Immediate complications after medical compared with surgical termination of pregnancy, *Obstet Gynecol*, 2009, vol. 114 (pg. 795-804)

[Google Scholar](#) [Crossref](#) [PubMed](#).

[Article Navigation](#)

[Google Scholar](#) [Crossref](#) [PubMed](#).

Parazzini F, Cipriani S, Chiaffarino F, Sandretti F, Bortolus R, Chiantera V. Induced abortion and risk of small-for-gestational-age birth, *Br J Obstet Gynaecol*, 2007, vol. 114 (pg. 1414-1418)

[Google Scholar](#) [Crossref](#)

Raatikainen K, Heiskanen N, Heinonen S. Induced abortion: not a risk factor for pregnancy outcome, but a challenge for health counselling, *Ann Epidemiol*, 2006, vol. 16 (pg. 587-592)

[Google Scholar](#) [Crossref](#) [PubMed](#).

Reime B, Schücking BA, Wenzlaff P. Reproductive outcomes in adolescents who had a previous birth or an induced abortion compared to adolescents' first pregnancies, *BMC Pregnancy Childbirth*, 2008, vol. 8 [10.1186/1471-2393-8-4](#).

[Crossref](#)

Shah PS, Zao J. Induced termination of pregnancy and low birth weight and preterm birth: a systematic review and meta-analysis, *Br J Obstet Gynaecol*, 2009, vol. 116 (pg. 1425-1442)

[Google Scholar](#) [Crossref](#)

Swingle HM, Colaizy TT, Zimmerman MB, Morriss FH. Abortion and the risk of subsequent preterm birth: a systematic review with meta-analyses, *J Reprod Med*, 2009, vol. 54 (pg. 95-108)

[Google Scholar](#) [PubMed](#).

THL (National Institute for Health and Welfare)The Medical Birth Register, 2012 <http://www.thl.fi/en/statistics/parturients>.

THL (National Institute for Health and Welfare)The Abortion Register, 2012 <http://www.thl.fi/en/statistics/abortions>.

van Oppenraaij RH, Jauniaux E, Christiansen OB, Horcajadas JA, Farquharson RG, Exalto N. ESHRE Special Interest Group for Early Pregnancy (SIGEP)Predicting adverse obstetric outcome after early pregnancy events and complications: a review, *Hum Reprod Update*, 2009, vol. 15 (pg. 409-421)

[Skip to Main Content](#)
[Google Scholar](#) [Crossref](#) [PubMed](#).

Virk J, Zhang J, Olsen J. Medical abortion and the risk of subsequent adverse pregnancy

[Article Navigation](#)

Voigt M, Henrich W, Zygmunt M, Friese K, Straube S, Briese V. Is induced abortion a risk factor in subsequent pregnancy?, *J Perinat Med*, 2009, vol. 37 (pg. 144-149)

[Google Scholar](#) [Crossref](#) [PubMed](#).

Yimin C, Wei Y, Weidong C, Xianmi W, Junqing W, Lin L. Mifepristone-induced abortion and birth weight in the first subsequent pregnancy, *Int J Gynecol Obstet*, 2003, vol. 84 (pg. 229-235)

[Google Scholar](#) [Crossref](#)

Zhou W, Sørensen HT, Olsen J. Induced abortion and low birthweight in the following pregnancy, *Int J Epid*, 2000, vol. 29 (pg. 100-106)

[Google Scholar](#) [Crossref](#)

© The Author 2012. Published by Oxford University Press on behalf of the European Society of Human Reproduction and Embryology. All rights reserved. For Permissions, please email:

journals.permissions@oup.com

Comments

1 Comment

Birth Outcomes After Induced Abortion

26 September 2012 | Colleen M Krajewski (with Colleen M. Krajewski, Roxanne Jamshidi, Anne E. Burke, Jennifer A. Robinson, Michelle C. Fox)

MD, MPH, Division of Family Planning, Johns Hopkins Bayview Medical Center

Sir,

We read with great interest the recently published nationwide register-based study of first births in Finland, which concluded that sexuality education should contain information on the potential hazards of repeat induced abortion (IA), including very preterm birth and low birthweight in subsequent pregnancies. The authors adjusted for 7 confounders, and found a significant association of several adverse pregnancy outcomes when comparing women with 0 vs 3 IA's. With the exception of very preterm birth, no dose- response relationship was demonstrated.

Sk

We found it interesting that a similar study in 2006 by Raatikainen et al using data from a representative region of Finland found no association between IA and birth outcomes. One explanation may be that

Article Navigation

- two major factors related to both spontaneous and indicated preterm birth. A Cochrane review of risk factors for preterm birth identified many of the same factors as important in risk scoring systems. We are concerned that important potential confounders such as these were not adjusted for in the current study.

The authors rightly noted that observational studies can only report associations, not causality and we wonder if the association between multiple IA's and preterm births persists once medical comorbidities and other factors are controlled for. The population-based database with which the authors have the privilege of working may contain valuable, yet untapped information that may shed further light on this interesting and important question.

Thank you,

Colleen M. Krajewski, Roxanne Jamshidi, Anne E. Burke, Jennifer A. Robinson, Michelle C. Fox

Division of Family Planning, Johns Hopkins Bayview Medical Center, Johns Hopkins Medical Institutions, Baltimore MD

Conflict of Interest:

Drs. Burke, Fox, and Jamshidi provide training for Nexplanon insertion. Dr. Krajewski (corresponding author) has no competing interests.

Submitted on 26/09/2012 8:00 PM GMT



View Metrics

Email alerts

[Article activity alert](#)

[Advance article alerts](#)

[New issue alert](#)

[Skip to Main Content](#)

[Receive exclusive offers and updates
from Oxford Academic](#)

Article Navigation

Factors affecting low birthweight after assisted reproduction technology: difference between transfer of fresh and cryopreserved embryos suggests an adverse effect of oocyte collection

Maternal age and risk of low birth weight and premature birth in children conceived through medically assisted reproduction. Evidence from Finnish population registers

What is the most relevant standard of success in assisted reproduction? Challenges in measuring and reporting success rates for assisted reproductive technology treatments: What is optimal?

Dilatation and curettage increases the risk of subsequent preterm birth: a systematic review and meta-analysis

Related articles in

Web of Science

Google Scholar

Related articles in PubMed

Preterm birth and the future risk of orthopedic fracture.

Junctional Zone Endometrium Morphological Characteristics and Functionality: Review of the Literature.

Stillbirth Among Arab Women in Canada, 1981-2015.

EPIDEMIOLOGIC CHARACTERISTICS AND RISK FACTORS FOR CONGENITAL HYPOTHYROIDISM FROM 2009 TO 2018 IN XIAMEN, CHINA.

[Skip to Main Content](#)

[Article Navigation](#)[Web of Science \(24\)](#)[Google Scholar](#)[Crossref](#)**Latest** | **Most Read** | **Most Cited**

Temporal trends in age at menarche and age at menopause: a population study of 312 656 women in Norway

Decidual NK cell-derived conditioned medium from miscarriages affects endometrial stromal cell decidualisation: endocannabinoid anandamide and tumour necrosis factor- α crosstalk

Exome sequencing reveals novel causes as well as new candidate genes for human globozoospermia

Type 1 diabetes in children born after assisted reproductive technology: a register-based national cohort study

Maternal age and risk of low birth weight and premature birth in children conceived through medically assisted reproduction. Evidence from Finnish population registers

[About Human Reproduction](#)[Editorial Board](#)[Policies](#)[Author Guidelines](#)[Contact ESHRE
Purchase](#)[Recommend to your Library](#)[Facebook](#)[Twitter](#)[YouTube](#)[LinkedIn](#)[Flickr](#)

Advertising and Corporate Services

Journals Career Network

Online ISSN 1460-2350

Print ISSN 0268-1161

Copyright © 2020 European Society of Human Reproduction and Embryology

[About Us](#)

[Contact Us](#)

[Careers](#)

[Help](#)

[Access & Purchase](#)

[Rights & Permissions](#)

[Open Access](#)

Resources

[Authors](#)

[Librarians](#)

[Societies](#)

[Sponsors & Advertisers](#)

[Press & Media](#)

[Agents](#)

Connect

[Join Our Mailing List](#)

[OUPblog](#)

[Twitter](#)

[Facebook](#)

[YouTube](#)

[Tumblr](#)

Explore

[Shop OUP Academic](#)

[Oxford Dictionaries](#)

[Oxford Index](#)

[Epigeum](#)

[OUP Worldwide](#)

[University of Oxford](#)

Oxford University Press is a department of the University of Oxford. It furthers the University's objective of excellence in research, scholarship, and education by publishing worldwide

OXFORD
UNIVERSITY PRESS

Copyright © 2020 Oxford University Press

[Cookie Policy](#)

[Privacy Policy](#)

[Legal Notice](#)

[Site Map](#)

[Accessibility](#)

[Get Adobe Reader](#)

[Article Navigation](#)