

See all >
[38 Citations](#)

See all >
[10 References](#)

[Download citation](#)

[Share](#) 

[Request full-text](#)

Buccal misoprostol as cervical preparation for second trimester pregnancy termination

Article in [Contraception](#) 65(6):415-8 · July 2002

with 56 Reads 

DOI: [10.1016/S0010-7824\(02\)00306-2](https://doi.org/10.1016/S0010-7824(02)00306-2) · Source:

[PubMed](#)

[Cite this publication](#)



Catherine S Todd
ORCID iD: [32.09](#) · Columbia University



Maria Soler



Laura Castleman



M Katherine Rogers



Paul D Blumenthal

[Hide](#)

Abstract

The objective of the study was to assess the efficacy of cervical preparation with misoprostol or laminaria suction abortion up to 18 weeks gestation. The study was conducted among a consecutive case series of patients presenting for dilatation and evacuation at a single center. Cervical preparation was effected by either 600 mg buccal misoprostol (n = 32) or laminaria (n = 78). Dilatation and evacuation was then performed using vacuum aspiration, with fetal parts or placenta removed as necessary with appropriate forceps. The principle outcome of interest was procedure time. Complete abortion was effected in 100% of the cases. In all cases, the cervix was adequately prepared to allow either the introduction of a size 14-mm suction cannula or was easily dilated to this diameter. Group size was sufficient for detection of a 20% difference in mean procedure time with 80% power (calculated with two separate standard deviations for procedure time). Procedure times were not significantly different when misoprostol was used compared to laminaria for cervical preparation. It is concluded that the use of misoprostol to provide cervical preparation in second trimester abortion procedures appeared effective and offers an alternative to laminaria. Buccal misoprostol should be more widely investigated for this purpose.

Do you want to **read the rest** of this article?

[Request full-text](#)



Citations (38)

References (10)

... The surgeon then administered a paracervical block with lidocaine and dilated the cervix using Hegar dilators. The uterus was evacuated according to the protocol outlined by Todd et al. [17, 18] After the abortion, women recovered at the clinic for approximately 2 h. One of the authors conducted extensive, informal, qualitative interviews with patients after the abortion in order to learn about women's perspectives on their care. ...

... To our knowledge, this is the first time MVA has been used as part of the D&E procedure in a low-resource setting such as Vietnam. We found two published retrospective studies in which buccal misoprostol was used for cervical preparation prior to D&E [18, 19]. Our data demonstrate that buccal misoprostol followed by MVA and forceps can be used in Vietnam to perform D&E up to 18 weeks of pregnancy. ...

Introduction of the dilation and evacuation procedure for second-trimester abortion in...

[Article](#)

Sep 2006 · [CONTRACEPTION](#)

Laura Castleman · Paul D Blumenthal ·  Oanh Khuat ·  Alyson Hyman

[View](#) [Show abstract](#)

... D&E can be performed safely under local anesthesia (paracervical block) with intravenous conscious sedation and analgesia. The most important requirements for safe D&E are the specialty training, skill and experience of the surgeon [18]. It is suggested that during forceps extraction the physician keep one hand on the fundus as a splint to reduce the risk of perforation. ...

... The fetal tissue should not be removed forcibly through the cervix, as bone spicules may lacerate the cervix. Crushing and rotating techniques lessen cervical trauma [18]. After forceps extraction, the uterine cavity should be gently explored with a large curette to ensure complete evacuation. ...

Maternal Klippel–Trenaunay Syndrome with Complete Hydrotidiform Mole and Coexistent

[Article](#)

Mar 2016

● Ahmed Samy El-agwany

[View Show abstract](#)

... The disadvantage includes the sometimes irritating, chalky taste. There are few clinical studies on buccal use of misoprostol including one for second-trimester abortion [11] and one for term labor induction [12]. There are several clinical studies on sublingual misoprostol for surgical abortion [13,14], missed abortion [15], term labor induction [16] and medical abortion [17]. ...

Comparison of misoprostol plasma concentrations following buccal and sublingual...

[Article](#)

Feb 2005 · CONTRACEPTION

Eric Schaff · ● Stephen L Fielding · Robert DiCenzo

[View Show abstract](#)

... No studies have been performed that address whether antibiotic administration at the time of dilator insertion would confer additional benefit. With use of misoprostol for cervical preparation prior to D&E, the risk of infection appears to be low^{48 49 50}. Two studies report no or few complications with misoprostol for cervical preparation but do not specifically report the number of observed infections [51,52]. ...

Prevention of infection after induced abortion: release date October 2010: SFP guideline 20102

[Article](#)

Apr 2011

● Sharon L. Achilles · Matthew F Reeves

[View](#)

... ate. Buccal misoprostol is becoming common

when used to prepare the cervix prior to surgical abortions and has also been used for labor induction. The only two studies comparing vaginal to buccal misoprostol used different doses: 200 µg buccal was compared to 50 µg vaginal to induce labor and there were no significant differences in efficacy [6,7]. [8]. At a National Abortion Federation meeting in April 2002, an informal survey of the several hundred abortion providers in the audience indicated that about one third were using buccal misoprostol. Pharmacokinetic studies have shown that sublingual misoprostol provides the greatest area under the curve of serum levels and so it is likely ...

Comparing vaginal and buccal misoprostol when used after methotrexate for early abortion

[Article](#)

Dec 2004 · [CONTRACEPTION](#)

● Ellen R Wiebe · Konia Trouton

[View Show abstract](#)

... Independent t test, values are mean circumference or french of the largest dilator that could be passed without resistance (SD). Todd et al 16 found that cervical preparation (defined as ease of dilation to 14 mm) and procedure time were similar in women randomized to receive either 600 mg buccal misoprostol or laminaria before surgical abortion between 14 and 18 weeks. No studies have established whether misoprostol alone will result in satisfactory cervical preparation in pregnancies beyond 18 weeks. ...

Cervical preparation using laminaria with adjunctive buccal misoprostol before second-...

[Article](#)

Mar 2006 · [Am J Obstet Gynecol](#)

● Alison Edelman · ● John Buckmaster · ● Martha F Goetsch · ● Jeffrey Jensen

[View Show abstract](#)

Unnatural Death of Foetus: A Forensic Study

[Book](#)

[Full-text available](#)

Jun 2019

● Ibtisam Sodani

[View](#)

Prostaglandins for induction of second-trimester termination and intrauterine death

[Article](#)

Nov 2003 · [BEST PRACT RES CL OB](#)

Suk Wai Ngai · Oi Shan Tang · ● Pak Chung Ho

[View Show abstract](#)

Prostaglandins for first-trimester termination

[Article](#)

Nov 2003 · [BEST PRACT RES CL OB](#)

● Philip Darney · ● Karen R Meckstroth

[View Show abstract](#)

Methods for Induced Abortion

[Article](#)

Jul 2004 · [OBSTET GYNECOL](#)

Phillip G. Stubblefield · ● Sacheen Carr Ellis · ● Lynn Borgatta

[View Show abstract](#)

Buccal or sublingual misoprostol for cervical ripening and labour induction

[Article](#)

Feb 2004 · [COCHRANE DB SYST REV](#)

Gibson Muzonzini · G Justus Hofmeyr

[View Show abstract](#)

Misoprostol Compared With Laminaria Before Early Second-Trimester Surgical Abortion: A...

[Article](#)

Sep 2005 · [OBSTET GYNECOL](#)

● Philip Darney · ● Amy Whitaker · ● Alisa B Goldberg · Eleanor A Drey

[View Show abstract](#)

Methods for Induced Abortion

[Article](#)

Feb 2005 · OBSTET GYNECOL

● Joseph R Siebert · Raj P Kapur · ● Robert Resta · ● David Luthy

[View Show abstract](#)

Uses of Misoprostol in Obstetrics and Gynecology

[Article](#)

[Full-text available](#)

Jul 2009

Rebecca Allen · ● Barbara M O'Brien

[View Show abstract](#)

Randomized trial of mifepristone and buccal or vaginal misoprostol for abortion through 56 day...

[Article](#)

Dec 2005 · CONTRACEPTION

Tamer Middleton · Eric Schaff · ● Stephen L Fielding · ● Beverly Winikoff

[View Show abstract](#)

Adequacy and safety of buccal misoprostol for cervical preparation prior to termination of...

[Article](#)

May 2006 · CONTRACEPTION

● Ashlesha Patel · Elizabeth Talmont · Johanna Morfesis · Vanessa Cullins

[View Show abstract](#)

Misoprostol administered by epithelial routes - Drug absorption and uterine response

[Article](#)

Sep 2006 · OBSTET GYNECOL

● Philip Darney · ● Karen R Meckstroth · ● Amy Whitaker · ● Suzanne M Bertisch

[View Show abstract](#)

A comparison of the efficacy of sublingual and oral misoprostol 400 microgram in the...

[Article](#)[Full-text available](#)Nov 2006 · [J Med Assoc Thai](#)Orada Patamasingh Na Ayudhaya · Y Herabutya · ●
Boonsri Chanrachakul · P O-Prasertsawat[View Show abstract](#)

Cervical preparation for second-trimester surgical abortion prior to 20 weeks of gestation[Article](#)Dec 2007 · [CONTRACEPTION](#)

● M. Fox · Jennifer L Hayes

[View Show abstract](#)

Recent advances in second-trimester abortion: an evidence-based review[Article](#)May 2009 · [Am J Obstet Gynecol](#)

Cassing Hammond

[View Show abstract](#)

Randomized trial of buccal versus vaginal misoprostol for induction of second trimester...[Article](#)

May 2010

● Sacheen Carr Ellis · ● Nathalie Kapp · Olivera Vragpvoc · Lynn Borgata

[View Show abstract](#)

Issues in second trimester induced abortion (medical/surgical methods)[Article](#)

Mar 2010

● Pak Chung Ho · ● Vivian CY Lee · Ernest Hung Yu Ng

[View Show abstract](#)

Laminaria vs. vaginal misoprostol for cervical preparation before second-trimester surgical...[Article](#)

Jan 2015

● Ron Sagiv · ● Yossi Mizrahi · Hagit Glickman ·
Abraham Golan

[View Show abstract](#)

Elective Pregnancy Termination

[Chapter](#)

May 2018

Cassing Hammond · Sheila Mody

[View Show abstract](#)

Synthetic Osmotic Dilators with Adjunctive Misoprostol for Same-day Dilation and...

[Article](#)

May 2016

Christy M. Boraas · ● Sharon L. Achilles · Miriam
Cremer · ● Beatrice A Chen

[View Show abstract](#)

Schwangerschaftsabbruch

[Chapter](#)

Jan 2006

H. Schneider · F. K. Beller

[View](#)

Induced Abortion Guidelines

[Article](#)

Nov 2006

Victoria Jane Davis

[View Show abstract](#)

Options for second-trimester termination

[Article](#)

Jan 2013

R. Perry · B. Harwood

[View Show abstract](#)

Misoprostol 1 to 3 h preprocedure vs. overnight

osmotic dilators prior to early second trimester

[Article](#)

Apr 2015

Shanthy Ramesh ·  Alicia Roston ·  Lindsay Zimmerman ·  Julie Chor

[View Show abstract](#)

Prevalence and risk factors of inadequate cervical dilation following laminaria insertion in second-...

[Article](#)

Jan 2015

 Ido Ben-Ami · Sharon Stern ·  Zvi Vaknin · Reuvit Halperin

[View Show abstract](#)

No. 360-Induced Abortion: Surgical Abortion and Second Trimester Medical Methods

[Article](#)

Jun 2018

Dustin Costescu ·  Edith Raphaëlle Guilbert

[View Show abstract](#)

Same-Day Synthetic Osmotic Dilators Compared With Overnight Laminaria Before Abortion at 14-...

[Article](#)

Jan 2014 · [OBSTET GYNECOL](#)

Eleanor A Drey ·  Sara J Newmann ·  Abby Sokoloff · Mithu Tharyil

[View Show abstract](#)

Directive clinique sur l'interruption volontaire de grossesse

[Article](#)

Nov 2006

Victoria Jane Davis · AUTEUR PRINCIPAL

[View Show abstract](#)

Cervical preparation for second trimester surgical abortion

[Chapter](#)

Jul 2008

● Sara J Newmann · ● Andrea Endres · ● Justin T
Diedrich · ● Jody Steinauer

[View](#)

Surgical Abortion in the Second Trimester

[Article](#)

May 2008

● Patricia A Lohr

[View Show abstract](#)

Cervical preparation for surgical abortion between 12 and 18 weeks of gestation using vaginal...

[Article](#)

Jun 2011

Leah C Wilson · ● Leslie A Meyn · Mitchell D Creinin

[View Show abstract](#)

Cervical preparation for second trimester dilation and evacuation

[Article](#)

Aug 2010 · COCHRANE DB SYST REV

Eleanor A Drey · ● Sara J Newmann · ● Andrea
Endres · ● Justin T Diedrich

[View Show abstract](#)

Second-trimester pregnancy termination with 600-??g vs. 400-??g vaginal misoprostol and...

[Article](#)

Feb 2008 · CONTRACEPTION

Josep L Carbonell · Miguel A Torres · Raúl Reyes ·
Carlos Sánchez

[View Show abstract](#)

Recommended publications

[Discover more publications, questions and projects in Misoprostol](#)

Article

Buccal misoprostol for cervical ripening prior to first trimester abortion

February 2010 · Contraception

● Ashlesha Patel · ● Hemang B. Panchal · ● Emily M. Godfrey

Cervical priming prior to uterine suction evacuation softens the cervix and lessens the force needed for dilation, thereby potentially reducing the probability of procedural complications. The use of buccal misoprostol has been shown to be an adequate cervical primer in second trimester surgical procedures, but its use in first trimester aspiration procedures is not well documented. Our objective ... [\[Show full abstract\]](#)

[Read more](#)[Discover more](#)

Last Updated: 27 Aug 2019

**Company**[About us](#)[News](#)[Careers](#)**Support**[Help](#)[Center](#)**Business solutions**[Advertising](#)[Recruiting](#)