

Search Our Scientific Publications & Authors

Search Publications & Authors

Publications ▼

Q SEARCH

FIND PUBLICATIONS BY CATEGORY +

Translate page: ▼

Immediate placement of intrauterine devices after first and second trimester pregnancy termination. (https://www.pubfacts.com/detail/21134501/Immediate-placement-of-intrauterine-devices-after-first-and-second-trimester-pregnancy-termination)

AUTHORS:

- Michelle C Fox (https://www.pubfacts.com/author/Michelle+C+Fox) Julia Oat-Judge (https://www.pubfacts.com/author/Julia+Oat-Judge)
- Kathryn Severson (https://www.pubfacts.com/author/Kathryn+Severson) Roxanne M Jamshidi (https://www.pubfacts.com/author/Roxanne+M+Jamshidi)
- Rameet H Singh (https://www.pubfacts.com/author/Rameet+H+Singh)
- Raegan McDonald-Mosley (https://www.pubfacts.com/author/Raegan+McDonald-Mosley) Anne E Burke (https://www.pubfacts.com/author/Anne+E+Burke)

Contraception 2011 Jan 11;83(1):34-40. Epub 2010 Aug 11.

🕒 January 2011

Department of Gynecology and Obstetrics, Johns Hopkins School of Medicine, Baltimore, MD 21287, USA.

📖 26 Reads

Background: We reviewed our experience with intrauterine device (IUD) placement after surgical abortion up to 20 weeks' gestation.

Study Design: Women presenting for elective abortion between January 2004 and March 2009 who requested an IUD were included in this retrospective review.

Results: Of 308 women requesting postabortion IUD placement, 221 (72%) planned insertion at the time of abortion (immediate group) and 87 (28%) planned insertion at their postoperative visit (interval group). IUDs were placed in 96% of the immediate group and in 23% of the interval group (212/221 vs. 20/87; p<.0001). Failure to return for placement was the most common reason for noninsertion in the interval group (60/87=69%). Follow-up information was obtained for 56% of patients and was documented a median of 137 days postabortion (range 3-1594 days). There was no difference in complication rates between groups. Expulsion rates were 3% and 0% in the immediate and interval groups, respectively (6/212 vs. 0/20; p=.4). Considering only those with documented follow-up after immediate insertion (119), there was a nonsignificant trend towards increased expulsion with placement after second vs. first trimester abortion (4/54=7% vs. 2/65=2%; p=.3). When analyzing the 172 subjects with documented follow-up, those planning immediate insertion were more likely to have an IUD in situ at the last contact than those planning later insertion (84/124=68% vs. 20/48=42%; p=.002).

(https://www.facebook.com/pubfacts) u=https://www.pubfacts.com/detail/21134501/Immediate-placement-of-intrauterine-devices-after-first-and-second-trimester-pregnancy-termination?text=Full+Article+on+PDF&mini=true&url=https://www.pubfacts.com/detail/21134501/Immediate-placement-of-intrauterine-devices-after-first-and-second-trimester-pregnancy-termination

Conclusion: Immediate postabortion IUD insertion is safe and effective. Given the low rate of return for interval insertion, immediate placement may be preferable.

Download full-text PDF

Source

<http://dx.doi.org/10.1016/j.contraception.2010.06.018> (<http://dx.doi.org/10.1016/j.contraception.2010.06.018>)

DOI Listing

PUBLICATION ANALYSIS

Top Keywords

- iud placement 8
- planned insertion 8
- interval group 8
- 2004 march 4
- january 2004 4
- 28% planned 4
- 2009 requested 4
- requested iud 4
- march 2009 4
- abortion january 4
- women presenting 4
- gestationstudy design 4
- postoperative visit 4
- presenting elective 4
- group 28% 4
- elective abortion 4

4
insertion postoperative
4
iud included
4
requesting postabortion
4
insertion time
4

Altmetric Statistics



Show full details (http://www.altmetric.com/details.php?citation_id=530005)

- 1 Total Shares (http://www.altmetric.com/details.php?citation_id=530005/).
- 1 Research Highlights (http://www.altmetric.com/details.php?citation_id=530005/research-highlights).
- 1 Citations (http://www.altmetric.com/details.php?citation_id=530005/).

MEMBER ARTICLES

Plasmodium vivax chloroquine resistance links to pvcr1 transcription in a genetic cross.



Dr. Juliana Martha Sa
Staff Scientist
National Institutes of Health
MD | United States

(<https://www.pubfacts.com/detail/31541097/Plasmodium-vivax-chloroquine-resistance-links-to-pvcr1-transcription-in-a-genetic-cross>)

The application of ceragenins to orthopedic surgery and medicine.



Dr Aiken Dao
Doctoral Researcher
Children's Hospital at Westmead
Westmead, NSW | Australia

(<https://www.pubfacts.com/detail/31994754/The-application-of-ceragenins-to-orthopedic-surgery-and-medicine>)

Clinical Aspects and Evolution under Current Treatments on Pulmonary Alveolar Phospholipoproteinosis Patients.



Dr. Petru Emil Muntean

Medic specialist pneumolog

Spitalul Judetean de Urgenta Arges, Dispensarul de Pneumoftiziologie Pitesti

Pitesti, Arges | Romania

<https://www.pubfacts.com/detail/31237860/Clinical-Aspects-and-Evolution-under-Current-Treatments-on-Pulmonary-Alveolar-Phospholipoproteinosis>

SIMILAR PUBLICATIONS



FACTS (<https://www.pubfacts.com/>)

© 2020 PubFacts.



About PubFacts (<https://www.pubfacts.com/about>)

Advertise (<https://www.pubfacts.com/advertise>)

Privacy Policy (<https://www.pubfacts.com/privacypolicy>)

Sitemap (<https://www.pubfacts.com/sitemap>)