

All Content ▾

Search [Advanced Search](#)

< Previous Article

1988 Volume 1, Issue 3, Pages 181–184

Next Article >

Access this article on
[ScienceDirect](#)

Medical treatment of urethral prolapse in the premenarchal female

[Lane J. Mercer](#), M.D. , [Christine M. Mueller](#), M.D., [Samir N. Hajj](#), M.D.

Section of Gynecology Department of Obstetrics and Gynecology, University of Chicago, Chicago, Illinois

PlumX Metrics

DOI: [https://doi.org/10.1016/S0932-8610\(88\)80058-6](https://doi.org/10.1016/S0932-8610(88)80058-6)

Abstract References

Abstract

Urethral prolapse, the circumferential protrusion of urethral mucosa through the urethral meatus, often presents in the premenarchal female as a large, friable mass. Some traditional therapies have included excision or cauterization. A series of 13 premenarchal females presented with urethral prolapse and were treated medically at the Chicago Lying-in Hospital. All patients had resolution of signs and symptoms within 4 weeks of initiation of therapy. No relapses or recurrences were noted. Medical therapy of urethral prolapse in the premenarchal female is a safe and successful alternative to surgery.

Key Words:

[Urethral prolapse](#), [Premenarchal female](#), [Medical management](#)

To access this article, please choose from the options below

Log In

[Login to existing account](#)

[Forgot password?](#)

Register

[Create a new account](#)

Purchase access to this article

- [\\$35.95 USD | PDF Download and 24 Hours Online Access](#)

Claim Access

If you are a current subscriber with Society Membership or an Account Number, [claim your access now](#).

Subscribe to this title

[Purchase a subscription](#) to gain access to this and all other articles in this journal.

Institutional Access

[Visit ScienceDirect](#) to see if you have access via your institution.

Article Tools

[PDF \(838 KB\)](#)

[Email Article](#)

[Add to My Reading List](#)

[Export Citation](#)

[Create Citation Alert](#)

[Cited by in Scopus \(0\)](#)

[Request Permissions](#)

[Order Reprints](#)

(100 minimum order)

Related Articles

A Review of Expedited Partner Therapy for the Management of Sexually Transmitted Infections in Adolescents

Kathryn E. Gannon-Loew, Cynthia Holland-Hall, Andrea E. Bonny

Vaginal dilator therapy: further suggestions for providers

Federica Facchin, Elisa Restelli, Paolo Vercellini

Vaginal Dilator Therapy: A Guide for Providers for Assessing Readiness and Supporting Patients Through the Process Successfully

Anne-Marie Amies Oelschlager, Katherine Debiec

Patient Delivered Expedited Partner Therapy for Chlamydia Among Female Adolescents Using

*This paper was presented in part at the Second Annual Meeting of the North American Society for Pediatric and Adolescent Gynecology held in Cleveland, OH Sept. 1987

© 1988 Springer-Verlag New York Inc. Published by Elsevier Inc. All rights reserved.

School Based Health Centers: Is It Feasible?

Susan H. Vacca, Melanie A. Gold, Elizabeth L. Salsgiver, Jeannette Rodriguez, Natalie Neu

Treatment Modalities in Adolescents Who Present with Heavy Menstrual Bleeding

Tasneem S. Alaqzam, Angela C. Stanley, Pippa M. Simpson, Veronica H. Flood, Seema Menon

[View All](#)

< Previous Article

1988 Volume 1, Issue 3, Pages 181–184

Next Article >

ELSEVIER Copyright © 2020 [Elsevier](#) Inc. All rights reserved. | [Privacy Policy](#) | [Terms & Conditions](#) | [Use of Cookies](#) | [About Us](#) | [Help & Contact](#) | [Accessibility](#)
The content on this site is intended for health professionals.

We use cookies to help provide and enhance our service and tailor content and ads. By continuing you agree to the [use of cookies](#).

Advertisements on this site do not constitute a guarantee or endorsement by the journal, Association, or publisher of the quality or value of such product or of the claims made for it by its manufacturer.

