
Email This		
Your Name:		
Your Email:		
Send me a copy		
Recipient's Email:		
Limit to one email		
Subject:		
Optional Message		
Limit 200 characters		
		//

Send



Abstracts Originally Presented at the National Abortion Federation



Search	
No Results	

Search No Results



Friday, April 17, 2020 News & Perspective Drugs & Diseases CME & Education Academy Video

MedGenMed Ob/Gyn & Women's Health > Meeting Proceedings

Scientific Paper Abstracts Originally Presented at the National Abortion Federation's Annual Meeting; April 23, 2007; Boston, Massachusetts

0 Read Comments

IN THIS ARTICLE

Serious Adverse Events Associated with Use of Misoprostol for Cervical Ripening in Early Second-Trimester Abortions (12-16 Weeks)

Mary Gatter, MD; Deb Nucatola, MD; Vallorie Saulsberry, MD, MPH

Introduction and Purpose/Objective(s): Our objective was to report on our experience with serious adverse events, particularly perforations, in a series of more than 6,000 cases of early second trimester abortions (12W0D to 16W0D) using 90 minutes of pre-op cervical preperation with 400 mcs of vaginal or buccal misoprostol. While there are a number of studies looking at misoprostol for cervical preperation in the first trimester, there are fewer studies examining

To continue reading this article ...

Log In

or

Register for Free

Experience Using a Metronidazole and Azithromycin Protocol for Prophylaxis at Time of Surgical Abortion in More Than 50,000 Women

Serious Adverse Events Associated with Use of Misoprostol for Cervical Ripening in Early Second-Trimester Abortions (12-16 Weeks)

The Medical Students for Choice Reproductive Health Externship Program: Impact on Medical Students' Knowledge, Attitudes, and Intention to Provide Abortions

Emotional Prophylaxis for Abortion Patients: A Pilot Study

Mifepristone Versus Laminaria: A Randomized Controlled Trial of Cervical Ripening in Midtrimester Induction

Obesity and Abortion

Misoprostol for Treatment of Incomplete Abortion in Madagascar and Moldova

Non-vaginal Routes of Misoprostol Administration for Pregnancy Termination up to 63 Days' LMP

Risk Factors for Repeat Elective Abortion

Can Women Accurately Assess the Success of Medication Abortion Based on Symptoms Alone?

Health Workers and Managers' Perspectives on Implementing Comprehensive Abortion Care in Ghana

What to Read Next on Medscape

Recommended Reading

2001/viewarticle/924583 JOURNAL ARTICLE

The Risks of Proton Pump Inhibitors During Pregnancy

2002259346-overview DISEASES & CONDITIONS Psychosocial and Environmental Pregnancy Risks

2001/viewarticle/928763 National Watchman Registry Reports Impressive Procedural Safety



2001/viewarticle/896156 JOURNAL ARTICLE Impact of TSH During the First Trimester of Pregnancy on Obstetric and Foetal Complications



1. Topol: US Betrays Healthcare Workers in Coronavirus Disaster

2. COVID-19: Could Hydroxychloroquine Really Be An Answer?

3. Toilets May Pose Risk for Spreading COVID-19

4. Coronavirus on Fabric: What You Should Know

5. Independent Nurse Practitioners and Physician Assistants: A Doc's View

View More



b:recommendationscuratedHasData : false



FIND US ON



ABOUT

- About Medscape
- **Privacy Policy**
- **Editorial Policy**
- Cookies
- Terms of Use
- Advertising Policy
- Help Center

MEMBERSHIP

Become a Member

Email Newsletters

Manage My Account

APPS

Medscape

CME & Education

WEBMD NETWORK

WebMD

MedicineNet

eMedicineHealth

RxList

Eropooio

WebMD Corporate

EDITIONS English Deutsch Español

Português

All material on this website is protected by copyright, Copyright © 1994-2020 by WebMD LLC. This website also contains material copyrighted by 3rd parties.

	Close